

Toxoplasma Retinitis: A Picture-Perfect Presentation

Shivani Chaturvedi*, Sandeep Saxena

Department of Ophthalmology, King George's Medical University, Lucknow, Uttar Pradesh, India.

Abstract

Toxoplasma Retinitis: A picture-perfect presentation where SD-OCT serves as a crucial supplemental imaging tool for both diagnosing and monitoring patients with ocular toxoplasmosis.

Keywords: Ocular toxoplasmosis, Spectral domain optical coherence tomography, Chorioretinitis.

INTRODUCTION

Toxoplasma gondii is an obligate, single-cell protozoan parasite that resides within host cells is responsible for ocular toxoplasmosis, an infection that primarily affects the retina and choroid. This condition ranks as the foremost cause of posterior uveitis globally and is a frequent contributor to vision impairment due to intraocular infections. *T. gondii* is highly prevalent in nature, with an estimated one billion people worldwide being infected by this parasite.

Risk factors for *T. gondii* infection include exposure to environments where the infectious organism is prevalent, particularly those frequented by cats. Additionally, individuals face an elevated risk of infection if they consume raw or undercooked meat containing *Toxoplasma* oocysts. Pregnant women infected with the parasite face a heightened risk of transmitting the infection to their unborn child, resulting in severe fatal complications, including retinal infections, congenital deformities, and mortality.

Ocular Toxoplasmosis

Toxoplasma infection leads to necrotizing chorioretinitis. This chorioretinal area with indistinct margins, accompanied by a focal vitreous infiltrate, is often described as “a headlight in the fog”. Additionally, localized lobular perivasculitis can occur around the arterioles in the vicinity of the active lesion, known as Kyrieleis plaques.¹ However, atypical lesions may present as extensive areas of retinal necrosis or retinochoroiditis without adjacent preexisting pigmented retinal scars, and they can affect both eyes. These unusual

lesions are more commonly seen in elderly individuals and individuals with underlying immunodeficiency caused by various factors.² Unconventional clinical presentations can be unfamiliar to healthcare providers, leading to delays in both diagnosis and treatment.

Visual impairment can result from a macular lesion, as seen in Fig. 1, whereas lesions in the peripheral retina often cause vision loss due to severe vitreous inflammation. Although less common, optic nerve involvement can lead to significant visual field defects and loss of color vision.³ Ocular toxoplasmosis may also give rise to additional manifestations, including retinal neovascularisation, retinal detachment and optic neuritis.

The polymerase chain reaction (PCR) method can be valuable in confirming the diagnosis, especially in immunocompromised individuals who exhibit extensive, atypical retinitis foci. On the other hand, assessing local antibody production may be a more suitable approach for accurately diagnosing immunocompetent individuals with smaller retinitis foci.⁴

Address for correspondence: Shivani Chaturvedi

Department of Ophthalmology, King George's Medical University, Lucknow, India.

E-mail: shivani.c4u@gmail.com

© UPJO, 2024 Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by-nc-sa/4.0/>.




Figure 1: Fundus Photo showing toxoplasma lesion over macula (A) Right Eye (B) Left Eye

Spectral Domain Optical Coherence Tomography in Ocular Toxoplasmosis

Spectral-domain optical coherence tomography (SD-OCT) has become a widely accepted technique for visualizing the posterior eye's structural details, offering an impressive resolution. SD-OCT imaging revealed that toxoplasma retinochoroidal lesions and scars exhibit a complex nature. In the acute phase, they are characterized by thickening and disorganization of the retinal reflective layers across their whole thickness and the underlying choroid.⁵ A thickened hyperreflective retina, accompanied by choroidal thickening and hyperreflectivity, hyperreflective dots in the vitreous,




Figure 2: SD-OCT showing toxoplasma lesion over macula (A) Right Eye (B) Left Eye




Figure 3: SD-OCT- 3D Visualisation of a macular cube (Right Eye)

and thickening of the posterior hyaloid are also seen (Figs 2 and 3). Adjacent to the lesion, there was frequently observed disorganization of retinal layers and interrupting of the ellipsoid zone (EZ). As time progressed, the retina underwent thinning and diminished hyperreflectivity, but it did not revert to a normal appearance. Additionally, there was thickening of the retinal pigment epithelium coupled with choroidal hyperreflectivity, or alternatively, epithelial atrophy with choroidal hyperreflectivity.⁶

As scar formation progresses, varying degrees of thinning become apparent, often accompanied by irregularities in the outer retinal layers.⁷

SD-OCT reveals discernible changes at the macula, with lesions characterized by hyperreflective dots at the vitreoretinal interface, full-thickness retinal hyperreflectivity, and choroidal thickening beneath the submacular region.^{8,9}

Crater like defects over macula pathognomonic of loss of larger areas associated with toxoplasmosis.

CONCLUSION

SD-OCT serves as a crucial supplemental imaging tool for both diagnosing and monitoring patients with ocular toxoplasmosis.

REFERENCES

1. Stokkermans TJ, Havens SJ. Toxoplasma Retinochoroiditis. [Updated 2023 Jun 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK493182/>
2. Fardeau C, Romand S, Rao NA, Cassoux N, Bettembourg O, Thulliez P, et al. Diagnosis of toxoplasmic retinochoroiditis with atypical clinical features. *Am J Ophthalmol.* 2002;134:196–203.
3. Ozgonul C, Besirli CG. Recent Developments in the Diagnosis

- and Treatment of Ocular Toxoplasmosis. *Ophthalmic Res.* 2017;57(1):1-12. doi: 10.1159/000449169. Epub 2016 Oct 11. PMID: 27723657.
4. Moshfeghi DM, Dodds EM, Couto CA, Santos CI, Nicholson DH, Lowder CY, et al. Diagnostic approaches to severe, atypical toxoplasmosis mimicking acute retinal necrosis. *Ophthalmology.* 2004;111:716–25.
 5. Oréfice JL, Costa RA, Scott IU, Calucci D, Oréfice F; Grupo Mineiro de Pesquisa em Doenças Oculares Inflamatórias (MINAS). Spectral optical coherence tomography findings in patients with ocular toxoplasmosis and active satellite lesions (MINAS Report 1). *Acta Ophthalmol.* 2013 Feb;91(1):e41-7. doi: 10.1111/j.1755-3768.2012.02531.x. Epub 2012 Sep 12. PMID: 22970804.
 6. Park JH, Lee SY, Lee EK. Morphological characteristics of ocular toxoplasmosis and its regression pattern on swept-source optical coherence tomography angiography: a case report. *BMC Ophthalmol.* 2019 Sep 5;19(1):199. doi: 10.1186/s12886-019-1209-8. PMID: 31488090; PMCID: PMC6729000.
 7. Goldenberg D, Goldstein M, Loewenstein A, Habot-Wilner Z, et al. Graefes Arch Clin Exp Ophthalmol. 2013 Aug;251(8):2037-45. Vitreal, retinal, and choroidal findings in active and scarred toxoplasmosis lesions: a prospective study by spectral-domain optical coherence tomography.
 8. Oliver GF, Ferreira LB, Vieira BR, Arruda S, Araújo M, Carr JM, Smith JR, Furtado JM, et al. Posterior segment findings by spectral-domain optical coherence tomography and clinical associations in active toxoplasmic retinochoroiditis. *Sci Rep.* 2022 Jan 21;12(1):1156. doi: 10
 9. Ammar F, Mahjoub A, Ben Abdesslam N, Knani L, Ghorbel M, Mahjoub H. Spectral optical coherence tomography findings in patients with ocular toxoplasmosis. *Ann Med Surg (Lond).* 2020 Apr 24;54:125-128. doi: 10.1016/j.amsu.2020.04.008. eCollection 2020 Jun.